

# Global System for Mobile communications (GSM) Electromagnetic Waves affect the Activity, Morphology, and Structure of Skeletal Muscles in Adult Male Rats

Wiam Ramadan, Hassan Khachfe, Lina Sabra, Khalil Abou Saleh, Mehnieh Baccouri, Wissam H. Joumaa

Department of Biological and Chemical Sciences  
Lebanese International University  
Nabatieh, Lebanon  
wiam.ramadan@liu.edu.lb, hassan.khachfe@liu.edu.lb

Environmental Physio-Toxicity Laboratory (R G017  
EPhyTox)  
Doctoral School for Science and Technology, Lebanese  
University  
Nabatieh, Lebanon  
lsabra@ul.edu.lb, khalil\_as@hotmail.com,  
mbaccouri@ul.edu.lb, wjoumaa@ul.edu.lb

Eric Esteve, Mohammed Moustafa ElSayed, Lina Ismail

HP2 Laboratory, INSERM U1042, Jean Roget Institute  
Joseph Fourier University,  
Faculty of Medicine and Pharmacy  
Grenoble, France  
eric.esteve@ujf-grenoble.fr

Molecular Biology and Genetics Laboratory  
Beirut Arab University, Faculty of Sciences  
Debbiyeh, Lebanon  
mohamed.moustafa@bau.edu.lb,  
linaismaill@hotmail.com

**Abstract**—The use of cellular technology is overwhelming our lives these days. Global System for Mobile communications (GSM) waves -the basis of cellular technology - are high frequency, high energy electromagnetic waves that may pose as a threat to man. The current work studies the effect of such waves on two types of skeletal muscles, the slow and fast twitching muscles in adult male rats. The activity, morphology, and structure of the affected muscles are studied and analyzed against control. Experiments evaluate changes in body weight, muscle mass, water and protein content, total RNA concentration, testosterone level, and Myosin Heavy Chain (MHC) isoforms expression. Our results show that in both muscles, there are changes in the distribution of muscles proteins and in the percentage of MHC isoforms suggesting that the GSM antenna relay affects the plasticity of skeletal muscle fiber by transforming slow type to faster one.

**Keywords**-GSM electromagnetic waves; skeletal muscles; proteins content; testosterone level; MHC isoforms.

## I. INTRODUCTION

The widespread use of mobile phones has been going sky-high over the past decade and the applications offered by mobile phone technology have become an essential part of personnel, business and social life. We have recently reported that the structure, the morphology and the activity of 2 types of striated muscles in rats have been altered by the exposition to electromagnetic waves emitted from mobile phones devices [1]. In fact, mammalian striated muscle myofibrils are composed of repeating units called sarcomeres that are arranged in series. Sarcomeres in turn are composed of contractile filaments termed myofilaments that are of two major types, actin (thin filament) and myosin (thick filament), which interact together to generate force

and contraction. These myofilaments are large polymers of noncovalently associated contractile proteins, actin and myosin, which comprise 70% of myofibrillar proteins in skeletal muscles [2]. The isomers of Myosin Heavy Chains (MHC) are often used to distinguish the types of skeletal muscle fibers: slow-twitch – or type I – muscle fibers, where MHC I isoform is abundantly expressed; and fast-twitch – or type II – muscle fibers (types IIa, IIb, and IIc/d), where MHC IIa, IIb, IIc/d predominate, respectively [3-5]. Slow-twitch fibers are adapted for continuous activity, and they are rich in myoglobin and oxidative enzymes. A typical example is the soleus muscle. Fast-twitch fibers are adapted for rapid activity, and they produce energy through glycolytic metabolism. A typical example is the *extensor digitorum longus (edl)* muscle [6].

A remarkable characteristic of striated muscles is plasticity. This term refers to the ability of these muscles to remodel and thus change their contractile and metabolic makeup, and – hence – their type from slow to fast, or vice versa, in response to specific environmental challenges, such as exercise, temperature, or gravitational loading, or internal challenges such as nutritional conditions as well as neuronal, mechanical, metabolic or hormonal stimuli [7]. This may be attributed to a reversible change in the muscle gene expression that leads to reversible structural and functional modifications [8].

One of the important challenges that have developed in the last decade and is thought to have an effect on health is the population exposure to electromagnetic waves, particularly the Global System of Mobile (GSM) communication signals. These signals are emitted from diverse sources particularly from cell phones and base station antennas. Also, they may come from industrial processes, where workers in broadcasting, transport, and

communication industries are highly exposed. They are also emitted from medical devices like electrosurgical devices and diagnosis equipment. Numerous studies have been conducted to measure, document, and archive the amount of Radio Frequency (RF) energy emitted from GSM sources in residential areas [9]. Thus, concerns from the risk of GSM signals on health arise from long term exposure, as well as from the cumulative effect of these waves. Researchers have been seeking options to minimize additional radiation exposures for the population and reduce the potential risk for harmful exposures.

Tyagi et al. studied the effect of mobile phone radiation on brain activity. They comparatively analyzed the effect of more than one type of radiations, and concluded that their effects on brain activity cannot be attributed to chance [10]. Tomruk et al. reported the substantial hepatic oxidative DNA and lipid damage on pregnant, non-pregnant, and newly-born rabbits [11]. Several studies reported the possibility that radio-frequency electromagnetic fields (EMF) used in cellular technology might influence DNA integrity of male germ cells as well as sperm motility. There was evidence that GSM radiation could induce, in vitro, the activation of stress conditions and response in human sperm cells [12]. The major mechanism by which such waves can induce an effect on biological systems is the thermal mechanism by which EMF at high intensities can increase the tissue or body temperatures above the normal value. Non-thermal mechanisms are under wide investigation in recent studies [13-16].

To understand the effects of EMF radiations, it would be beneficial to appreciate the nature and mode of action of this type of energy. EMF radiation is a form of potential energy exhibiting wavelike behavior as it travels in the space.

EMF radiation has both electric and magnetic field components that oscillate – in phase – perpendicular to each other and orthogonal to the direction of energy propagation [16]. Such radiations can be classified as ionizing and non-ionizing radiation, based on capability to ionize atoms and/or molecules, and to break chemical bonds. The non-ionizing type is associated with potential non-thermal hazards. The short high energy or the chronic low energy exposure of biological tissues to EMF radiations has been shown to change the functional activities of cells, resulting finally in some diseases [10].

Few studies have investigated the effect of electromagnetic waves on skeletal muscles. In fact, Radicheva *et al.* (in 2002) has shown that a 2.45 GHz microwave field could possess a stimulating effect on muscle fiber activity, which is in part due to its specific non-thermal properties [18]. Moreover, our previous study has shown that one hour of exposure to electromagnetic field at 900 MHz modulated by human voice could have an effect on the excitation-contraction coupling mechanism of mammalian fast-twitch skeletal muscles [19]. However, no

study to date has investigated the effect of electromagnetic waves emitted by GSM relay antenna on muscle composition. Consequently, this study is designed to investigate the effect of 25V/m of electromagnetic waves emitted by GSM relay antenna on animal body weight, muscle mass, proteins and water content, total RNA expression, serum testosterone level and myosin heavy chain isoforms expression in the two types of skeletal muscle fibers, slow and fast-twitches.

Kaasik et al. have shown that MHC I isoform relative content in human muscle was 2.6 times higher than in horse and 6.3 times higher than in rat muscle. This may be related to the differences in endurance capacity of human, horse, and rat muscle [20]. Indeed, these authors have shown that the main difference between the distribution of myosin light chain (MLC) isoforms in human, horse, and rat skeletal muscle is the relatively low level of regulatory MLC isoforms in human skeletal muscle. MLC isoforms distribution in skeletal muscle has been shown to be related to the physiological role and adaptational capacity of muscle to everyday motor activity [21]. Despite the relative proportions of each fiber type vary between homologous muscles of different species [22-23], and in the absence of human data, research with experimental animals is the most reliable means of detecting important toxic properties of chemical substances and for estimating risks to human and environmental health. Thus, we decided to use as animal model, the rat, in order to study the effects of GSM electromagnetic waves on skeletal muscles. The exposures received by animals can be compared to those received by humans in order to interpret test results and predict risk. This in turn helps regulatory agencies to prioritize funding for environmental cleanup.

This paper studies the effect of GSM waves on skeletal muscles in rats. A background of the study is given in Section I. The materials and methods used in the study are mentioned in Section II. The results are presented in Section III and discussed in Section IV.

## II. MATERIALS AND METHODS

### A. Experimental Design

All procedures in this study were performed in accordance with the stipulations of the Helsinki Declarations and with the current Lebanese laws for animal experimentation. Twenty adult Sprague-Dawley male rats with an average weight of  $190 \pm 5$  g were divided equally into 2 groups. One group was subjected for 6 weeks to whole continuous (24 hours/day) body exposure to EMW (900 MHz,  $E_{\text{eff}} = 25\text{V/m}$ ) (Fig. 1). The other group was considered as control and maintained in the same environmental conditions under the turned off antenna. Both exposed and control animals were housed in a temperature-controlled room (22°C) on a 12:12-h light-dark cycle. They were daily supplied with the same kind of food and water.




Fig. 1. GSM electromagnetic Waves exposure system

### B. Dissection

After the exposure period, the rats were gently sacrificed and trunk blood was collected. Soleus and *edl* muscles were rapidly excised from the hind limbs of each rat. The muscles are weighted and preserved at  $-80^{\circ}\text{C}$  for later analysis.

### C. Total RNA extraction

Total RNA was extracted from muscle samples using **RiboZol™ RNA Extraction Reagent** from AMRESCO, according to the venter's instructions (American Research Products, 30175 Solon Industrial Parkway, Solon, OH 44139-9827 USA).

### D. Serum testosterone level determination

Collected blood was centrifuged at 3500 rpm for 5 minutes. Serum of each rat was preserved at  $-20^{\circ}\text{C}$ . Serum testosterone, insulin and cortisol levels, in control and exposed groups, were measured by Enzyme-Linked ImmunoSorbent Assay (ELISA) technique based on the principle of competitive binding, according to instructions supplied by the venter.

### E. Proteins dosage

Protein dosage was performed according to Bradford Technique [24]. Pieces of frozen muscles were mechanically disrupted and spliced in 5 volumes of washing buffer containing 20 mM NaCl, 1 mM EGTA (pH 6.4), and 5 mM  $\text{PO}_4$ . After 5 minutes of centrifugation at a high speed (12000 rpm), the supernatant is collected and the quantity of the protein is determined with the Bradford method (Bio-Rad, Hercules, CA), where the results were expressed as a ratio of milligrams of proteins to 100 milligrams of muscles. The pellet was then washed with 3 volumes of extraction buffer containing 5 mM EGTA, 1mM

dithiothreitol (pH 8.5), and 100 mM sodium pyrophosphate, and incubated in cold overnight. The next day, the mixture was centrifuged at 12000 rpm for 10 minutes and the supernatant – which contained the protein myosin – was collected and the amount of myofibrillar proteins was determined. Small volumes (50 $\mu\text{L}$ ) of the supernatant were diluted twice with glycerol and stored at  $-20^{\circ}\text{C}$  for electrophoresis.

### F. SDS-PAGE electrophoretic separation of Myosin Heavy Chain isoforms

To analyze the content of MHC I, MHC IIa, IIb, IIc/x isoforms in the extracts, we used simple vertical migration of SDS-PAGE electrophoretic separation. The separating gel was prepared from 99.5% glycerol, 30% acrylamide, 0.6% bis acrylamide, 1.5 M Tris (pH 8.8), 1 M glycine, 10% SDS, 10% ammonium persulfate, and TEMED. The stacking gel was prepared from 99.5% glycerol, 30% acrylamide, 0.6% bis acrylamide, 0.5 M Tris (pH 6.8), 10% SDS, 0.1 M EDTA (pH 7), 10% ammonium persulfate, and TEMED. For best quantification, 2-3  $\mu\text{g}$  of myosin were loaded in each well. Electrophoresis was performed using a Cleavage, Scientific ltd, system. Gels were run at constant voltage (70V) for 24 h and then stained with silver reagent that allowed the detection of the MHC bands corresponding to I, IIa, IIb, and IIc/x isoforms. The stained gels were scanned using a Canon digital imaging system and the density of bands was estimated using the UN-Scan-IT software [25].

### G. Statistical analysis

All values are expressed as means  $\pm$  SE for n observations. Data were analyzed by One-Way ANOVA

(StatView; Alsyd, Meylan, France) statistical test. A level of  $p < 0.05$  indicated statistical significance.

### III. RESULTS

#### A. Effect of GSM waves exposure on Body mass

As shown in Fig. 2, all animals steadily gained weight and there was no difference observed between the control and exposed animals after 6 weeks of GSM waves exposure (Control:  $283 \pm 8$  g; Exposed:  $295 \pm 7$  g,  $n=10$ )




Fig. 2. Effect of GSM electromagnetic waves exposition on Body mass. Each value displays mean  $\pm$ SE.

#### B. Effect of GSM waves exposure on muscles mass

Although body weight was not affected by the exposure, six weeks of exposure resulted in a significant decrease in *edl* mass by 16% (Control;  $133.56 \pm 3.69$  mg; Exposed:  $112.19 \pm 2.57$  mg,  $n=20$ ,  $p < 0.05$ ). However, no significant effect was observed in soleus muscle mass (Control;  $120.06 \pm 2.89$  mg; Exposed:  $117.49 \pm 3.11$  mg,  $n=20$ ) (Fig. 3).

Such a decrease in muscle mass observed in *edl* muscle could be related to modification in water content and/or in the proteins content. Consequently, the water content and the soluble and myofibrillar proteins content were estimated.

#### C. Effect of GSM waves exposure on muscles water content

In control group, water content expressed as percentage is estimated to  $37.5 \pm 0.7\%$  in soleus muscles and  $26.9 \pm 0.6\%$  in *edl* muscles. After 6 weeks of continuous electromagnetic waves exposure, and although no significant effect was observed in soleus muscle mass, an increase by 17% in percentage of water content was observed ( $43.9 \pm 1.2\%$ ,  $n=12$ ,  $p < 0.05$ ). However, no significant effect was observed in *edl* muscles ( $28.1 \pm 0.6\%$ ,  $n=12$ ) (Fig. 4).




Fig. 3. Effect of GSM waves exposure on muscles mass of the *edl* and soleus muscles. Each histogram displays mean  $\pm$ SE. \*  $p < 0.05$




Fig. 4. Effect of GSM electromagnetic waves exposition on muscles water content of the *edl* and soleus muscles. Each histogram displays mean  $\pm$ SE. \*  $p < 0.05$

#### D. Effect of GSM waves exposure on proteins content

In soleus control muscles, soluble and myofibrillar proteins content were  $2.71 \pm 0.13$  and  $3.42 \pm 0.26$  mg/g of muscle, respectively. The six weeks of continuous electromagnetic waves exposure induced an increase by 23% of soluble proteins ( $3.34 \pm 0.16$  mg/g,  $n=24$ ,  $p < 0.05$ ); however, a decrease by 32% of myofibrillar proteins was observed ( $2.33 \pm 0.29$  mg/g,  $n=24$ ,  $p < 0.05$ ) (Fig. 5).

In *edl* control muscles, soluble and myofibrillar proteins content were  $3.56 \pm 0.18$  and  $3.83 \pm 0.11$  mg/g of muscle, respectively. Six weeks of continuous GSM waves exposure induced a decrease by 28% and 24% of soluble and myofibrillar proteins, respectively (Soluble proteins content:  $2.78 \pm 0.19$  mg/g; Myofibrillar proteins content:  $2.91 \pm 0.22$  mg/g of muscle,  $n=24$ ,  $p < 0.05$ ) (see Fig. 5).

These modifications in proteins content in both soleus and *edl* muscles should be correlated to total RNA level expression.

**E. Effect of GSM waves exposure on total RNA**

In control condition, the total RNA values were  $755.21 \pm 16.72$   $\mu\text{g}/\mu\text{l}$  and  $548.32 \pm 14.98$   $\mu\text{g}/\mu\text{l}$  in *edl* and soleus muscles, respectively. After 6 weeks of GSM exposure, a decrease by 29% and 25% were shown in both *edl* and soleus muscles, respectively ( $n=10$ ;  $p < 0.05$ ) (Fig. 6).

This decrease in the amount of total RNA expression could be related to the modification in the serum testosterone level.




Fig. 5. Effect of GSM waves exposure on Myofibrillar proteins content and soluble proteins content on both soleus and *edl* muscles. Each histogram displays mean  $\pm$ SE. \*  $p < 0.05$




Fig. 6. Effect of GSM waves exposure on total RNA in both soleus and *edl* muscles. Each histogram displays mean  $\pm$ SE. \*  $p < 0.05$

**F. Effect of GSM waves exposure on serum testosterone, insulin and cortisol level**

In control condition, the Enzyme-Linked ImmunoSorbent Assay showed that serum testosterone level was  $82.12 \pm 4.39$  ng/ml. Six weeks of GSM waves exposure induced a decrease by 50% in serum testosterone level ( $40.91 \pm 5.71$  ng/ml,  $n=10$ ,  $p<0.05$ ) (Fig. 7).




Fig. 7. Effect of GSM waves exposure on testosterone levels in both control and exposed rats. Each histogram displays mean  $\pm$ SE. \*  $p < 0.05$

The serum insulin level shows a significant increase by 30% in rats exposed to the GSM (Fig. 8).




Fig. 8. Effect of GSM waves exposure on insulin levels in both control and exposed rats. Each histogram displays mean  $\pm$ SE. \*  $p < 0.05$

On the other hand, no significant difference was observed in the serum cortisol level after 6 weeks of GSM exposition (Fig. 9).

**G. Effect of GSM waves exposure on Myosin Heavy Chain Isoforms expression**

Separation and analysis of MHC isoforms by SDS-PAGE allowed the estimation of the density of the bands corresponding to each of the MHC isoforms (MHC I, II<sub>a</sub>, II<sub>b</sub>, and II<sub>x</sub>) using the UN-Scan-IT software. These isoforms are differentially expressed in the different muscle fiber types.

In control conditions, the *edl* muscle expresses  $34.6 \pm 0.2$  % of MHC II<sub>x</sub> and  $66.4 \pm 2.8$  % of MHC II<sub>b</sub>, while the soleus muscle expresses  $5.1 \pm 1.7$ % of MHC II<sub>a</sub> and  $94.6 \pm 1.5$ % of MHC I.

The 6 weeks of GSM waves exposure induced, in *edl* muscle, a significant increase by 63% in the expression of MHC II<sub>x</sub> isoforms ( $56.3 \pm 4.2$ %) and a significant decrease by 33% in MHC II<sub>b</sub> isoforms expression ( $44.9 \pm 3.4$ %,  $n=24$ ,  $p<0.05$ ).

Moreover, in soleus muscle, the exposure induced a significant increase in the expression of MHC II<sub>a</sub> isoforms ( $16.1 \pm 1.7$ %) with a significant decrease in the expression of MHC I isoforms ( $84.3 \pm 1.5$ %,  $n=24$ ,  $p<0.05$ ) (Fig. 10).




Fig. 9. Effect of GSM waves exposure on cortisol levels in both control and exposed rats. Each histogram displays mean  $\pm$ SE.




Fig. 10. Separation and analysis of MHC isoforms by SDS-PAGE in soleus and *edl* muscles in control rats and in rats exposed to 6 weeks of GSM waves. This figure show the bands corresponding to each of the MHC isoforms (MHC I, II<sub>a</sub>, II<sub>b</sub>, and II<sub>x/d</sub>)

**IV. DISCUSSION**

Many questions were raised about the possibility that exposure to electromagnetic fields emitted by mobile

phones or their base stations could affect the health of users. If there is a health impact, there will be a global impact because the number of active cell phones is estimated to reach 7.3 billion by 2014. For this, the World Health Organization (WHO) established the International EMF Project in 1996 to assess the science, and recommends research to fill gaps in the knowledge of risks arising from exposure to electromagnetic fields on health [8]. Safety is a valid concern for the users of wireless equipment, especially in regard to possible hazards caused by electromagnetic fields (EMFs). There has been increasing concern about the possible adverse health effects resulting from exposure to radiofrequency radiations (RFR), such as those from mobile communication devices, in general, and cellular phones, in particular. Mobile communication technology is achieved when signal is transferred via electromagnetic wave through radio frequency and microwave signals. This signal produces electromagnetic radiation in the form of thermal radiation that consists of harmful ionizing radiation and harmless non-ionizing radiation. When using mobile phone, electromagnetic wave is transferred to the body, potentially causing health problems, especially – for example – at the place near the ear skull region, where they are known to affect the neurons. Neurophysiological studies identified that radiations interfere with the electrical impulses that two neurons connect each other with. This can lead to adverse effects, from headache to blurred vision, to deafness and migraines. People using cell phones are susceptible to high blood pressure and other symptoms such as headaches and fatigue [26, 27]. There have been various studies into the connection between mobile phones and memory and the results were confusing and sometime contradictory [28-30].

Other studies have explored the possibility that radio-frequency electromagnetic fields (RF-EMF) employed by mobile phone technology could influence the genetic makeup and physiology of some specific cells, i.e., male germ cells, and sperm mobility [12]. It was shown that upon prolonged exposure to RF-EMF, sperm surface area and acrosomal region were significantly reduced. It was also shown that the primary function of sperm cells – binding to human oocytes – was significantly compromised.

Furthermore, the effect of RF-EMF radiations was studied vis-à-vis their induction of cellular stress response. This was evaluated by quantifying the expression and activity of heat shock proteins (Hsp), namely Hsp27 and Hsp70, but was shown not to have measurable differences from normal [12].

Skeletal muscle is the most abundant tissue in animals representing up to 50% of body mass in some athletic species such as dogs and horses. Muscle fibers are composed of myofibrils arranged in parallel, which constitute the major compartment in muscle cells, comprising from 73.2% of muscle fiber volume in horses to 83.3% of muscle fiber volume in goats. Also, total myofibrillar volume is directly proportional to muscle mass with a scaling factor of 0.98 [31].

Mammalian skeletal muscle is a highly heterogeneous tissue, able to adapt to environmental challenges to which it is subjected. In rats and mice, this process is governed by a set of mechanical, hormonal and nutritional signals [32-34]. Phenotypic plasticity of muscle tissue allows it to be modified in order to meet the specific demands faced by an animal during its life [18]. This critical property leads to a conversion of muscle fibers from slow to fast or vice versa [3, 35]. A wide range of contractile properties are mainly attributed to the diversity of the isoforms of MHC, which can exist in different muscle fibers. Four MHC isoforms (I, IIa, IIx and IIb), each encoded by a separate gene can be expressed in adult skeletal muscle. The intrinsic differences in the properties of the ATPase of MJHC isoforms led to the classification of fiber muscle as slow or fast fibers [33, 36]. Generally, the fast-type genes appear to be expressed at birth, while the slow-type genes are expressed in response to changes in activity during development [32].

This study aims to examine the effect of electromagnetic waves emitted from GSM antenna relay, at 900 MHz frequency, on body weight, muscle mass, protein content and the expression of the isoforms of the myosin heavy chain (MHC) in 2 types of muscle fiber types. One is slow oxidative fiber and the other is fast glycolytic fiber.

Actually, very few studies have examined the effect of electromagnetic waves on the skeletal muscle. In one of them, the study was conducted in vivo and showed that the field of microwave of 2.45 GHz has a stimulating effect on the activity of the muscle fibers, which is in part due to its non-thermal specific properties [18]. In another, the study was conducted in vitro, and showed that one hour of exposure to 900 MHz of electromagnetic field modulated by the human voice could have an effect on the mechanism of excitation-contraction coupling of skeletal muscle fast-twitch [19].

The results of our present work show that electromagnetic waves do not affect the body weight of rat males. In addition, the exposure period results in a decrease in absolute edl muscle mass, while that of soleus is kept unchanged. The myofibrillar protein content is directly proportional to muscle mass. A decrease in the mass of the edl was translated by decrease of both myofibrillar protein content and soluble protein. The increase in soluble protein content in soleus muscle was compensated by the decrease of myofibrillar proteins content resulting in the maintenance of muscle mass of soleus. This can be explained by the compensatory mechanisms at the translational and post-translational levels as suggested by Nikolova et al. [37]. In addition, synthetic modification in myofibrillar and soluble proteins in edl and soleus may be the consequence of the changes of level of transcription of genes encoding these proteins or due to perturbation in the stability of their corresponding mRNAs [35]. Therefore, the quantification of the mRNA encoding the soluble and myofibrillar and measurement of protein half-life is necessary to detect the level at which the change has led to modify the level of

myofibrillar proteins and other proteins expressed in these two types of skeletal muscles. Our results provide new information by showing a decrease in the total RNA values in both edl and soleus muscles after 6 weeks of exposure to GSM in rats. These results are very propitious and may explain some of disorders observed in humans exposed to electromagnetic waves. However, further studies on human are needed in order to confirm these possible effects.

On the other hand, the effect of electromagnetic waves is specific and can be attributed to a modification in the activity of different hormones. In 2010, Meo et al. [38] showed that exposure to radiation emitted by mobile phone for 60 minutes / day for a period of 3 months significantly decreased the level of serum testosterone in albino rats Wistar. In 2008, Al-Akhras showed that exposure for 6 weeks in a sinusoidal electromagnetic field of 50 Hz resulted in a significant reduction in levels of gonadotropins (FSH and LH) in female rats [39]. These results are similar to our results, where we showed a decrease in the level of testosterone after 6 weeks of exposure to GSM waves. This decrease can explain the decrease in total RNA values in both muscles since it is known that testosterone binds with androgen receptors inside the nucleus of different target cells and turn on the synthesis of mRNA, which are then translated into specified proteins. Thus, the decrease of the level of testosterone may explain the decrease of the RNA levels. On the other hand, there was no change in the level of cortisol in the rats exposed to GSM electromagnetic waves. This result suggests that the exposition to these radiations will not affect the hypothalamus-pituitary-adrenal axis that regulates the stress behavior. Thus, it is possible to propose that the radiations emitted from GSM will not induce stress in rats exposed to them. However, our results show an increase in the level of insulin in rats exposed to electromagnetic waves. This finding is similar to those shown by Meo and Rubeann [40] who have demonstrated that exposure to mobile phones causes hyperglycemia with a simultaneous and significant increase in serum insulin levels. Eventually, a compensatory rise in insulin level helps to control the hyperglycemia. Considering this result and its connection with mobile phone radiation, it is important to find out whether it is associated with B-cell dysfunction or insulin resistance.

If insulin resistance was developed, this can be linked to the change in the percentage of MHC isoforms observed in the present study. In fact, our results show an increase of the MHC IIa isoform, but a decrease in the MHC I isoform in the soleus muscle. In the edl muscle, a significant increase in MHC IIx and a significant decrease of MHC IIb isoforms were observed. These results suggest that the GSM antenna relay affects the plasticity of skeletal muscle fiber by transforming slow type to faster one. Thus, insulin resistance decreases the entry of glucose in muscle cells. These cells will switch from the slow aerobic type to the fast glycolytic one in order to accommodate the decrease in glucose uptake. These suggestions need to be established by

confirming the development of the insulin resistance and by studying the glucose uptake by muscles cells in rats exposed to mobile phone radiation.

#### ACKNOWLEDGMENTS

This work was supported by the National Council for Scientific Research (CNRS) in Lebanon, the Doctoral School of Science and Technology (EDST) at the Lebanese University, and the Francophone University Agency (AUF).

#### REFERENCES

- [1] W. Ramadan et al., Effect of GSM Electromagnetic Waves on the Activity, Morphology, and Structure of Skeletal Muscles, The Second International Conference on Global Health Challenges, IARIA Conference, ISBN: 978-1-61208-314-8, pp 17-22, Lisbon, Portugal, 17-21 November, 2013
- [2] M. J. Greenberg, J. D. Watt, M. Jones, K. Kazmierczak, D. Szczesna-Cordary, and J. R. Moore, Regulatory Light Chain Mutations Associated with Cardiomyopathy Affect Myosin Mechanics and Kinetics, *J Mol Cell Cardiol*, 2009, 46(1), pp. 108-115.
- [3] S. Ennion, J. Sant'ana Pereira, A. J. Sargeant, A. Young, and G. Goldspink, Characterization of human skeletal muscle fibers according to the myosin heavy chains they express, *J Muscle Res Cell Motil*, 1995, 16, pp. 35-43.
- [4] S.P. Fauteck and S.C. Kandarian, Sensitive detection of myosin heavy chain composition in skeletal muscle under different loading conditions, *Am J Physiol*, 1995, 268, pp. C419-424.
- [5] F. Chemello, C. Bean, P. Cancellara, P. Laveder, C. Reggiani, and G. Lanfranchi, Microgenomic Analysis in Skeletal Muscle: Expression Signatures of Individual Fast and Slow Myofibers, *PLoS One*, 2011, 6 (2), pp. 1-14.
- [6] S. Schiaffino and C. Reggiani, Myosin isoforms in mammalian skeletal muscle, *J Appl Physiol*, 1994, 77, pp. 493-501.
- [7] T. Soukup and I. Jirmanová, Regulation of myosin expression in developing and regenerating extrafusal and intrafusal muscle fibers with special emphasis on the role of thyroid hormones, *Physiol. Res.* (2000), 49, pp. 617-633.
- [8] T. Leeuw and Pette D, Coordinate change in the expression of troponin subunit and myosin heavy chain isoforms during fast -to-slow transition of low - frequency -stimulated rabbit muscle, *Eur J biochem* 1993, 213, pp. 1039-1046.
- [9] T. Haumann, U. Munzenberg, W. Maes, and P. Sierck, HF-Radiation levels of GSM cellular phone towers in residential areas, unpublished report.
- [10] A. Tyagi, M. Duhan, and D. Bhatia, Effect of mobile phone radiation on brain activity GSM VS CDMA, *IJSTM*, 2011, Vol. 2, Issue 2, pp. 1-5.
- [11] A. Tomruk, G. Guler, and A.S. Dincel, The influence of 1800 MHz GSM-like signals on hepatic oxidative DNA and lipid damage in nonpregnant, pregnant, and newly born rabbits, *Cell Biochem Biophys.*, 2010, 56(1), pp. 39-47
- [12] N. Falzone, C. Huyser, F. Fourie, T. Toivo, D. Leszczynski and D. Franken, In vitro effect of pulsed 900 MHz GSM radiation on mitochondrial membrane



- potential and motility of human spermatozoa, *Bioelectromagnetics*, 2008, 29(4): pp. 268-276
- [13] R.K. Adair, Biophysical limits on athermal effects of RF and microwave radiation, *Bioelectromagnetics*, 2003, 24(1), pp. 39-48.
- [14] A. Ahlbom, A. Green, L. Kheifets, D. Savitz, and A. Swerdlow, Epidemiology of Health Effects of Radiofrequency Exposure, *Environmental Health Perspectives*, 2004, 112(17), pp. 1741-1754.
- [15] M. Blettner and G. Berg, Are mobile phones harmful? *Acta Oncol*, 2000, 39(8), pp. 927-930.
- [16] C. Cinel, R. Russo, A. Boldini, and E. Fox, Exposure to Mobile Phone Electromagnetic Fields and Subjective Symptoms: A Double-Blind Study, *Psychosomatic Medicine*, 2008, 70, pp. 345-348.
- [17] R. Serway and J. Jewett, *Physics for Scientists and Engineers*, 2010, 9<sup>th</sup> ed. Brooks/Cole, Boston, USA.
- [18] N. Radicheva, K. Mileva, T. Vukova, B. Georgieva, and I. Kristev, Effect of microwave electromagnetic field on skeletal muscle fibre activity, *Arch Physiol Biochem*, 2002, 110(3), pp. 203-214.
- [19] W. Ramadan et al., Effect of electromagnetic field GSM on contractile responses and fast-twitch intact skeletal muscle fibers, *The FASEB Journal*, 2011, 25, 1051.15.
- [20] P. Kaasik, K. Leisson, R. Puhke, K. Alev, and T. Seene, Characteristics of myosin isoforms in mammalian skeletal muscle, *Advances in Biological Chemistry*, 2012, 2, pp. 77-83
- [21] K. Alev et al., Physiological role of myosin light and heavy chain isoforms in fast- and slow-twitch muscles: Effect of exercise, *Biology of Sport*, 2009, 26, pp. 215-234
- [22] M. A. Ariano, R. B. Armstrong, and V. R. Edgerton, Hindlimb muscle fiber populations of five mammals, *J Histochem Cytochem*, 1973, 21 (1), pp. 51-55.
- [23] I. Fuentes, A. R. Cobos, and L. A. G. Segade, Muscle fibre types and their distribution in the biceps and triceps brachii of the rat and rabbit, *J Anat*, 1998, 192 (2), pp 203-210.
- [24] R. Krauspe and A. Scheer, Coomassie brilliant blue G-250 dye-binding technique for determination of autolytic protein breakdown in *Euglena gracilis* and comparison to other methods of autolysis measurement, *Anal Biochem*, 1986, 153(2), pp. 242-250.
- [25] T.A. Kohn and K.H. Myburgh, Electrophoretic separation of human skeletal muscle myosin heavy chain isoforms: The importance of reducing agents, *J Physiol Sci*, 2006, 56 (5), pp. 355-360
- [26] S. Braune, A. Riedel, J. Schulte-Mönting, and J. Raczek, Influence of a radiofrequency electromagnetic field on cardiovascular and hormonal parameters of the autonomic nervous system in healthy individuals, *Radiat Res*, 2002, 158 (3), pp. 352-356.
- [27] S. Braune, C. Wrocklage, J. Raczek, T. Gailus, and C. H. Lücking, Resting blood pressure increase during exposure to a radio-frequency electromagnetic field, *Lancet*, 1998, 351 (9119), pp. 1857-1858.
- [28] A. Barth et al., A meta-analysis for neurobehavioural effects due to electromagnetic field exposure emitted by GSM mobile phones, , 2008, 65 (5), pp. 342-346.
- [29] M. S. Kwon and H. Hamalainen, Effects of mobile phone electromagnetic fields: critical evaluation of behavioral and neurophysiological studies, *Bioelectromagnetics*, 2011, 32 (4), pp. 253-272.
- [30] C. Wiholm et al., Mobile phone exposure and spatial memory, *Bioelectromagnetics*, 2009, 30 (1), pp. 59-65.
- [31] G. Goldspink, Changes in muscle mass and phenotype and the expression of autocrine and systemic growth factors by muscle in response to stretch and overload, *J. Anat*, 1999, 194, pp. 323-334
- [32] L. Larsson L, D. Biral, M. Campione, and S. Schiaffino, An age-related type IIb to IIx myosin heavy chain switching in rat skeletal muscle, *Acta Physiol Scand*, 1993, 147, pp. 227-234.
- [33] M. Nocella, B. Colombini, M.A. Bagni, J. Bruton, and G. Cecchi, Non-crossbridge calcium-dependent stiffness in slow and fast skeletal fibres from mouse muscle, *J Muscle Res Cell Motil*, 2012, 32(6), pp. 403-409.
- [34] S. Schiaffino and C. Reggiani, Molecular diversity of myofibrillar proteins: gene regulation and functional significance, *Physio Rev*, 76(2), pp. 371-423.
- [35] S.D. Harridge, Plasticity of human skeletal muscle: gene expression to in vivo function, *Exp Physiol*, 2007, 92(5), pp. 783-797.
- [36] C.E. Pandorf, W.H. Jiang, A.X. Qin, P.W. Bodell, K.M. Baldwin, and F. Haddad, Calcineurin plays a modulatory role in loading-induced regulation of type I myosin heavy chain gene expression in slow skeletal muscle, *Am J Physiol Regul Integr Comp Physiol*, 2009, 297(4), pp. R1037-1048.
- [37] T. Nikolova et al., Electromagnetic fields affect transcript levels of apoptosis-related genes in embryonic stem cell-derived neural progenitor cells. *The FASEB Journal*, 2005, 10.1096/fj.04-3549fje.1.
- [38] S.A. Meo, A. M. Al-Drees, S. Husain, M.M. Khan, and M.B. Imran, Effects of mobile phone radiation on serum testosterone in Wistar albino rats, *Saudi Med J*, 2010, 31(8), pp. 869-873.
- [39] M.A. Al-Akhras, Influence of 50 Hz magnetic field on sex hormones and body, uterine, and ovarian weights of adult female rats, *Electromagn Biol Med*, 2008 27(2), pp. 155-163.
- [40] S. A. Meo and K. Al Rubeaan, Effects of exposure to electromagnetic field radiation (EMFR) generated by activated mobile phones on fasting blood glucose, *Int J Occup Med Environ Health*, 2013, 26 (2), pp. 235-241.